

GRS 1915+105: the flux ratio of twin radio clouds as a measure of asymmetry between counter jets

A. M. Atoyan^{1,2} and F. A. Aharonian¹

¹ Max-Planck-Institut für Kernphysik, Postfach 10 39 80, 69029 Heidelberg, Germany

² Yerevan Physics Institute, Alikhanian Brothers 2, 375036 Yerevan, Armenia

Abstract

Resolution of both approaching and receding ejecta in the galactic microquasars makes possible to measure the flux ratio S_a/S_r of twin ejecta, which contains an important information about the nature of the jets. We show that the flux ratio $S_a/S_r = 8 \pm 1$ observed from GRS 1915+105 during the prominent March/April 1994 radio flare can be explained in terms of relativistic motion of discrete radio clouds, if one assumes that the twin ejecta are similar, but not completely identical, i.e. allowing for some asymmetry between the plasmoids in their speeds of propagation and/or luminosities. The recoil momentum due to asymmetrical ejection of the pair of plasmoids could be comparable with the momentum accumulated in the inner accretion disk. We suggest a possible explanation of the observed anticorrelation between the X-ray and radio flares as the result of drastic structural changes in the inner disk caused by production of powerful jets. The delay between the times of decline of X-rays and appearance of strong radio flares is explained by time needed for expansion of radio clouds to become optically thin.

Subject headings: stars: individual (GRS 1915+105) – stars: flare – galaxies: jets – radio continuum: general – accretion disks

1 Introduction

Recent discovery of the galactic superluminal jet sources GRS 1915+105 (Mirabel & Rodriguez 1994, hereafter MR94) and GRO J1655-40 (Tingay et al. 1995; Hjellming & Rupen 1995) provides unique possibility for deep study of the phenomenon of relativistic jets observed elsewhere in the Universe. These sources, called *microquasars*, represent, to a large extent, scaled down analogs of AGNs (Mirabel & Rodriguez 1995). Being, however, much closer to us than AGNs, the microquasars enable radio monitoring of *both* approaching and receding relativistic ejecta, and importantly, in short timescales.

GRS 1915+105 is the first source of apparent superluminal jets where the receding jet has been also detected, and time evolution of radio fluxes of both components of the prominent 19 March 1994 flare has been traced on timescales of days beyond 30 April (MR94). In particular, this has allowed Mirabel & Rodriguez to determine the real speed $\beta \approx 0.92$ and the angle of propagation $\theta \approx 70^\circ$ of the ejecta. At an estimated distance $D = 12.5 \pm 1.5$ kpc, the observed proper motion of radio sources correspond to apparent velocities (in units of speed of light) $v_a = 1.25 \pm 0.15$ and $v_r = 0.65 \pm 0.08$ for the approaching and receding ejecta, respectively (MR94).

Detection of pairs of jets makes possible to put forward an important question concerning the origin of the radio images observed. Namely, interpretation of the ratio of radio fluxes detected from the twin jets in GRS 1915+105 actually addresses an important issue, whether the moving radio images

are the radio ‘echos’ of relativistic shocks propagating in the jets, or they correspond to real motion of discrete plasmoids.

The ratio of flux densities, $S(\nu) \propto \nu^{-\alpha}$, measured at equal angular separations of the *identical* pair of jets from the core is

$$\frac{S_a}{S_r} = \left(\frac{1 + \beta \cos \theta}{1 - \beta \cos \theta} \right)^{k+\alpha}. \quad (1)$$

The index k is equal to $k = 3$, if the fluxes are produced in the moving *discrete* radio clouds, but $k = 2$ for the brightness ratio of continuous *stationary* jets (e.g. see Lind & Blandford 1985). The difference in the index k is due to relativistic volume contraction effect (see Bodo & Ghisellini 1995). For $\beta = 0.92$, $\theta = 70^\circ$, and measured radio spectral index $\alpha = 0.84 \pm 0.03$ (MR94), Eq.(1) results in the flux ratio $S_a/S_r \simeq 12$ to be expected from discrete sources, whereas the measured flux ratio $S_a/S_r = 8 \pm 1$ (MR94) is closer to $S_a/S_r \simeq 6$ which is formally expected in the case of stationary jets. Meanwhile, the apparent motion of discrete radio condensations does not support the interpretation of the observed flux ratio in terms of emission of stationary jets. A possible explanation of this discrepancy between the observed and expected (for discrete plasmoids) flux ratios of the jets in GRS 1915+105 was given by Bodo & Ghisellini (1995), who suggested that the real speed of radioemitting plasma (*fluid*) in the jets is different from the speed $\beta \approx 0.92$ of radio *patterns* attributed to propagation of shocks in the fluid.

In this paper we propose another interpretation of the measured flux ratio, in terms of speeds of the ejecta coinciding with the speeds $\beta \sim 0.9$ of radio patterns, assuming that the twin ejecta are *similar*, but *not* absolutely identical.

2 Pair of asymmetrical jets

When both components of relativistic jets can be detected, then in addition to the equation for the angular speed μ_a of the approaching jet, two more equations become available, namely, the one for the angular speed μ_r of the receding component, as well as the equation for the flux ratio of the two components. Allowing for the parameters of the counter ejecta to be different, the first two equations read:

$$\mu_a = \frac{\beta_a \sin \theta_a}{1 - \beta_a \cos \theta_a} \frac{c}{D}, \quad (2)$$

$$\mu_r = \frac{\beta_r \sin \theta_r}{1 + \beta_r \cos \theta_r} \frac{c}{D}, \quad (3)$$

where β_a and β_r are the speeds, θ_a and θ_r are the angles of propagation of approaching and receding ejecta. Note that for convenience in Eq.(3) we have substituted $\theta_r \rightarrow 180^\circ - \theta_r$.

The equation for the flux ratio S_a/S_r for an asymmetrical pair of ejecta is found, using the relation $S(\nu) = \delta^{3+\alpha} S'(\nu)$ (e.g., Lind & Blandford 1985) between the apparent and intrinsic energy fluxes of a radio cloud moving with Doppler factor $\delta = \sqrt{1 - \beta^2} / (1 - \beta \cos \theta)$. Then the flux ratio at equal *intrinsic times* is:

$$\frac{S_a}{S_r} = \left(\frac{\Gamma_r}{\Gamma_a} \right)^{3+\alpha} \left(\frac{1 + \beta_r \cos \theta_r}{1 - \beta_a \cos \theta_a} \right)^{3+\alpha} \frac{L'_a}{L'_r}, \quad (4)$$

where $\Gamma_{a,r} = (1 - \beta_{a,r}^2)^{-1/2}$ are the Lorentz factors of the bulk motion, and $L'_{a,r}$ are the intrinsic luminosities of the clouds. An asymmetry between the ejecta would imply at least one of the following inequalities: (a) the luminosities of the approaching and receding ejecta in their rest frame are different, i.e. $L'_a \neq L'_r$; (b) the ejecta propagate not strictly in the opposite directions, i.e. $\theta_a \neq \theta_r$; (c) the speeds of the ejecta are different, $\beta_a \neq \beta_r$. In principle, one may assume also that the spectra of the ejecta are described by different power-law indexes α . However, since the same $\alpha \approx 0.84$ was observed from both radio clouds of the 19 March 1994 flare, we do not consider here such a possibility.

Eq.(4) corresponds to the flux ratio of two radio clouds at equal *intrinsic times*, $t'_a = t'_r = t'$. Meanwhile, the directly measurable quantity is the flux ratio at equal *angular separations* $\phi_a = \phi_r = \phi$. The relation between these two quantities, $(S_a/S_r)_{t'}$ and $(S_a/S_r)_\phi$, can be found taking into account that in the observer's frame the ratio of the times corresponding to equal angular separation of the sources from the core is $(t_a/t_r)_\phi = \mu_r/\mu_a$, while the ratio of apparent times $t_{a,r} = t'/\delta_{a,r}$ corresponding to the same intrinsic time t' is $(t_a/t_r)_{t'} = \delta_r/\delta_a$. Using Eqs. (2) and (3), we find that $(t_a/t_r)_\phi = \lambda(t_a/t_r)_{t'}$ where

$$\lambda = \frac{\Gamma_r \beta_r \sin \theta_r}{\Gamma_a \beta_a \sin \theta_a} . \quad (5)$$

Approximating the time evolution (decline) of the flare as $S_{a,r} \propto t^{-p}$, as it was for the 19 March 1994 flare of GRS 1915+105 (with $p \simeq 1.3$, Mirabel & Rodriguez 1995), the relation between the flux ratios at equal intrinsic times and equal angular separations is found:

$$\left(\frac{S_a}{S_r}\right)_{t'} = \lambda^p \left(\frac{S_a}{S_r}\right)_\phi . \quad (6)$$

For GRS 1915+105 calculations below result in $\lambda \sim 0.85$, therefore the measured $(S_a/S_r)_\phi \sim 8$ corresponds to $(S_a/S_r)_{t'} \sim 6.5$. Further on we will omit subscript (t') , implying under (S_a/S_r) the flux ratio at equal intrinsic times, if not mentioned otherwise.

Returning now to Eqs. (2)–(4), note that for identical jets, when $\beta_a = \beta_r = \beta$, $\theta_a = \theta_r = \theta$, Eq.(5) results in $\lambda = 1$, therefore in this particular case the equal angular separations correspond to equal intrinsic times, as expected, so $S_a/S_r = (S_a/S_r)_\phi$. Taking into account also that $L'_a = L'_r$, Eq.(4) comes to Eq.(1) with $k = 3$. Then Eqs. (1)–(3) make formally a system of 3 equations for 3 variables: β , θ , D . However, these equations are *not* independent when the power-law index k is fixed, since they *predict* the ratio $S_a/S_r = (\mu_a/\mu_r)^{k+\alpha}$. Note that due to high value of the index $k + \alpha \simeq 4$, the predicted flux ratio is very sensitive to variations of the angular velocities $\mu_{a,r}$ of the twin ejecta. In particular, in the case of $k = 3$ and $\alpha = 0.84$, small error limits of the observed values of $\mu_a = 17 \pm 0.4$ mas/d (milliarcsec/day) and $\mu_r = 9.0 \pm 0.1$ mas/d (MR94) result in a rather broad range of the flux ratios predicted for identical plasmoids, $(S_a/S_r)_\phi = S_a/S_r = 13.1^{+1.8}_{-1.6}$, which is, nevertheless, beyond the observed $(S_a/S_r)_\phi = 8 \pm 1$.

However, this discrepancy disappears when an asymmetry between the ejecta is allowed. Consider separately each of the 3 basic types of asymmetries mentioned above.

A. $\beta_a = \beta_r = \beta$ and $\theta_a = \theta_r = \theta$, but $L'_a \neq L'_r$. In this case the solution to Eqs. (2) and (3) for GRS 1915+105 is given in MR94: $\beta \cos \theta = 0.323$, and for the distance $D = 12.5$ kpc the true speeds of the ejecta $\beta \approx 0.92$ and $\theta \approx 70^\circ$. Then from Eq.(4) follows that in order to explain the ratio $(S_a/S_r)_\phi \sim 8$, the ratio of intrinsic luminosities should be $L'_r/L'_a \sim 1.6$.

B. $\beta_a = \beta_r = \beta$ and $L'_a = L'_r$, but $\theta_a \neq \theta_r$. Introducing dimensionless parameters $v_a = \mu_a D/c$ and $v_r = \mu_r D/c$, which correspond to the apparent speeds (in units of speed of light) of the ejecta for a given distance D , the solution to Eqs. (2)–(4) can be written as:

$$\tan \theta_a = \frac{2(s-1)v_a}{(s-1)^2 + v_a^2(b^2 - 1)}, \quad (7)$$

$$\sin \theta_r = b \sin \theta_a, \quad (8)$$

$$\beta = \frac{v_a}{\sin \theta_a + v_a \cos \theta_a}, \quad (9)$$

where $s \equiv (S_a/S_r)^{1/(3+\alpha)}$, and $b \equiv sv_r/v_a = s\mu_r/\mu_a$. Eqs. (7)–(9) define θ_a , θ_r (in the range $\leq 180^\circ$) and β for arbitrary μ_a , μ_r and s (note that if $b > 1$, one should care for the condition $\sin \theta_r \leq 1$). However, starting from some distances these equations result in $\beta > 1$, which becomes formally possible after reduction of Γ_a and Γ_r in Eq.(4). For the values of μ_a and μ_r observed for the March 19 event, and $S_a/S_r \leq 7$, the solutions with $\beta < 1$ require $D < 10.7$ kpc. For example, assuming $D = 10$ kpc and $S_a/S_r = 7$, Eqs. (7)–(9) result in $\theta_a = 83.7^\circ$, $\theta_r = 57.7^\circ$ and $\beta = 0.92$. From Eq.(8) follows that symmetrical solutions, with $\theta_a = \theta_r$, exist only if $b = 1$, which is not the case in GRS 1915+105.

C. $L'_a = L'_r$ and $\theta_a = \theta_r = \theta$, but $\beta_a \neq \beta_r$. In this case, due to very strong dependence of the flux ratio S_a/S_r on the ratio Γ_a/Γ_r (see Eq.(4)), the discrepancy between the observed and ‘expected’ flux ratios by factor of (1.5 – 2) could be easily explained with only $\sim (10 - 20)\%$ difference in Lorentz factors of the oppositely moving plasmoids. Dividing Eq.(2) to Eq.(3), and substituting the resulting equation into Eq.(4), we find the relation between β_r and β_a , and afterwards Eqs. (2) and (3) can be resolved with respect to β_a and $\chi \equiv \beta_a \cos \theta$:

$$\chi = \frac{(1 - b^2)v_a^2 + s^2 - 1}{(1 - b^2)v_a^2 + s^2 + 2sb + 1}, \quad (10)$$

$$\beta_a = \sqrt{\chi^2 + v_a^2(1 - \chi)^2}, \quad (11)$$

$$\beta_r = \beta_a \frac{b}{\sqrt{1 + (b^2 - 1)\beta_a^2}}. \quad (12)$$

Parameters s and b are the same as in Eqs.(7)–(9). The angle θ is readily found after calculations of χ and β_a .

In Fig.1 we show the dependence of the speeds of propagation, β_a and β_r , of the approaching and receding plasmoids on the distance D , calculated from Eqs. (10)–(12) for $\mu_a = 17$ mas/d, $\mu_r = 9$ mas/d, and fixed flux ratio at equal intrinsic times $S_a/S_r = 7$. For the distance $D = 12.5$ kpc the speeds $\beta_a = 0.926$ and $\beta_r = 0.902$, and the angle $\theta = 70.2^\circ$ are found. From Eq.(6) follows that in this case the flux ratio $S_a/S_r = 7$ corresponds to $(S_a/S_r)_\phi \simeq 8.6$. Remarkably, the speeds of both ejecta are close to the value of $\beta = 0.92$ found in MR94 assuming $\beta_a = \beta_r$. In Fig.2 we show the speeds β_a and β_r and relevant Lorentz factors Γ_a and Γ_r of counter ejecta, calculated for the same $\mu_{a,r}$ and α as in Fig.1, but for different flux ratios in a broad range $5 \leq S_a/S_r \leq 15$. Note that *small* power-law index $1/(3 + \alpha) = 0.26$ in the parameter $s = (S_a/S_r)^{1/(3+\alpha)}$ significantly reduces the impact of uncertainties in the flux ratio S_a/S_r on calculated speeds of propagations of the ejecta.

3 Discussion

The key assumption of this paper consists in allowance for an asymmetry between the pair of plasmoids, which implies at least one of the enequalities: $L'_a \neq L'_r$, $\theta_a \neq \theta_r$, or $\beta_a \neq \beta_r$. Assuming that only the angles of propagation are different, $\theta_a \neq \theta_r$, one can explain the observed flux ratio $(S_a/S_r)_\phi = 8 \pm 1$ if GRS 1915+105 would be at a distance $D \simeq 10$ kpc, and requiring essential difference ($\geq 25^\circ$) between θ_a and θ_r . Note, however, that in this case the distance to GRS 1915+105 does not agree with estimated kinematic distance $D = 12.5 \pm 1.5$ kpc (MR94). Besides, the radio observations of the March 19 outburst clearly show the motion of two sources in strictly opposite directions on the sky, within uncertainties of only few degrees (MR94). Therefore the difference in the angles θ_a and θ_r could be only in the plane of view.

More probable seem to be realization of the other two types of asymmetries. If the asymmetry is attributed mainly to different intrinsic (i.e. in the rest frames of the clouds) luminosities $L'_a \neq L'_r$, then the receding cloud should be significantly ($\geq 50\%$) more luminous than the approaching one. In principle, such a difference in the luminosities of approaching and receding clouds could be attributed to different (by $\sim 30\%$) magnetic fields B , and/or to different amount of relativistic electrons in the ejecta. At last, the observed flux ratio can be explained with only a small difference in the speeds of propagation, $\beta_a \neq \beta_r$, corresponding to $\simeq 10\%$ difference between the Lorentz factors Γ_a and Γ_r of the bulk motion of the plasmoids. Due to very strong dependence of S_a/S_r on the ratio Γ_a/Γ_r , a small asymmetry in the speeds of propagation of the jets can be considered as the main reason for the discrepancy between the measured and ‘expected’ flux ratios in GRS 1915+105. Note, however, that there is no particular reason to believe that only the velocities of the ejecta are different, and it is quite possible that the intrinsic luminosities of the clouds can be also somewhat different, $L'_a \sim L'_r$ (but $\theta_a \approx \theta_r$). In that case Eqs. (10)–(12) can be used for calculations of the jet parameters, if one substitutes parameter $s = (S_a/S_r)^{1/(3+\alpha)}$ there with $s_* = (S_a L'_r/S_r L'_a)^{1/(3+\alpha)}$.

The asymmetry in the luminosities and/or speeds of propagation of the ejecta could be connected both with somewhat different conditions in the external medium at distances $\leq 10^{16} - 10^{17}$ cm from the core (reached by the ejecta in a few days) or with an asymmetry of the bidirectional ejection itself. In the absence of a firmly established theoretical model for development of the two-sided jets, there is no particular reason to eliminate any of these two options. For example, different densities of the external medium might result in different speeds of propagation, and/or different speeds of *expansion* of radio clouds leading to different magnetic fields in the clouds at equal intrinsic times. An asymmetry of the bidirectional ejection (which is obvious for GRO J1655-44, see Hjellming & Rupen 1995), e.g. $\sim 10\%$ difference in the masses or acceleration rates of the ejecta, could also result in different speeds of propagation and/or luminosities of the plasmoids.

Remarkably, intrinsic asymmetry of bidirectional ejection process could imply an interesting interpretation of the anticorrelation between the X-ray and radio flares observed from both microquasars GRS 1915+105 (Foster et al. 1996; Harmon et al. 1997) and GRO J1655-44 (Tingay et al. 1995; Harmon et al. 1995; Hjellming & Rupen 1995), which consists in appearance of strong radio outbursts generally after essential decline of the preceding X-ray flares, usually with some delay up to few days between these events. Tingay et al. (1995) and Meier (1996) interpreted this effect as an indication of

suppression of jet production mechanisms in the regime of supercritical accretion responsible for strong X-ray flares, while ejection would occur only after slowing down of the accretion rate to subcritical regime. Possible asymmetry of the jets allows us to suggest an alternative (in fact, to a large extent opposite) scenario where the jet production itself may result in temporary drop of the accretion from supercritical to subcritical regimes.

Indeed, internal energy in the radio clouds, and correspondingly, minimum kinetic energy of the ejection for strong outbursts in GRS 1915+105 is estimated as $W \sim 10^{43}$ erg (Liang & Li 1995; Meier 1996), and even more (MR94). Then the momentum transferred to relativistic ejecta can be estimated as $P_{a,r} \sim W/c \geq 3 \times 10^{32}$ g cm/s. Asymmetrical ejection generally would require, just from momentum conservation law, that significant *recoil* momentum $\Delta P = |\mathbf{P}_a - \mathbf{P}_r|$ is transferred to the core of ejection, which is the third object in the interacting system "two jets + core". Since most probable site responsible for relativistic ejection is the inner part of the accretion disk (e.g., Blandford & Payne 1982; Begelman et al. 1984), it is interesting to compare ΔP with the integrated value of the specific (per unit volume) momentum $\rho v = \rho |\mathbf{v}|$ of the gas orbiting in the inner accretion disk at radii $R' \leq R$: $P_d(R) = \int \rho v d^3 R'$ (here ρ is the mass density in the accretion flow). Using the Shakura & Sunyaev (1973) accretion disk model, this integrated 'absolute momentum' is estimated (in their *a*-region) as $P_d \simeq 4 \times 10^{21} \alpha_{SS}^{-1} \dot{m}^{-1} r^3$ g cm/s, where $\alpha_{SS} \leq 1$ is the viscosity parameter, and $r = R/R_g$ is the radius in the units of $R_g = 3(M/M_\odot)$ km. Then, even for $\alpha_{SS} \sim 0.01$, the black hole mass $m = M/M_\odot \sim 10$, and accretion rate about the critical, $\dot{m} \sim 1$, a small recoil momentum $\Delta P \sim 0.1 P_{a,r}$ being applied to the jet production region (approximately, perpendicular to the disk plane) would be enough to cause significant changes, or even destruction, of the inner disk up to radii $r_* \sim 100$ estimated from condition $\Delta P \sim P_d(r_*)$, and resulting in a temporary reduction/termination of the fuel supply into this region responsible for thermal X-rays.

Thus, in the framework of this scenario, the onset of subcritical/supercritical accretion would correspond to the active state, with X-ray flares which could proceed without significant ejection events and observable radio flares (see Foster et al. 1996; Tavani et al. 1996). Powerful ejection of radio emitting material may be accompanied by significant destruction of the inner accretion zone, leading to a strong decline of the X-ray fluxes simultaneously with production of relativistic ejecta. The ejection event, however, will be not accompanied by the *simultaneous* increase of the radio fluxes, since for an appreciable time the ejecta remain too compact to be transparent at the radio frequencies. Thus, one has to expect a delay, up to a few days, between the instant of ejection and the appearance of strong radio flares, which time is needed for expansion of the plasmoids, with $v_{\text{exp}} \sim 0.1 c$, to the radii $R_{\text{cl}} \geq 5 \times 10^{14}$ cm, as follows from the optical transparency of the radio plasmoids with respect to synchrotron self-absorption (Atoyan & Aharonian 1997). Note that the radio clouds, moving with the speeds $v \sim c$, become transparent at distances $R \sim 10 R_{\text{cl}} \geq 5 \times 10^{15}$ cm, i.e. far away from the binary system. Depending on the timescale needed for the accretion disk to recover after ejection event, the maximum of radio flares could appear just in the dips between X-ray flares (see Harmon et al. 1997). Rough estimate of the duration Δt of such dips as the dynamical time of accretion from radii r_* results in $\Delta t \propto r_*^{7/2}$, which could vary on time scales from minutes to days, depending on the spatial scales of the inner disk affected by ejection.

Acknowledgments We thank H.J.Völk for valuable discussions. AMA acknowledges the support of his work through the Verbundforschung Astronomie/Astrophysik of the German BMBF under the grant No. 05-2HD66A(7), and hospitality of the Max-Planck-Institute für Kernphysik during his visit to Heidelberg.

References

- Atoyan, A. M., & Aharonian, F. A. 1997, Proc. Workshop "Relativistic jets in AGN", Cracow, (in press)
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1984, Rev. Mod. Phys., 56, 255
- Blandford, R. D., & Payne, D. G. 1982, MNRAS, 199, 883
- Bodo, G., & Ghisellini, G. 1995, ApJ, 441, L69
- Foster, R. S., et al. 1996, ApJ, 467, L81
- Harmon, B. A., et al. 1995, Nature, 374, 704
- Harmon, B. A., et al. 1997, ApJ, 477, L85
- Hjellming, R. M., & Rupen, M. P. 1995, Nature, 375, 464
- Liang, E., & Li, H. 1995, A&A, 298, L45
- Lind, K. R., & Blandford, R. D. 1985, ApJ, 295, 358
- Meier, D. 1996, ApJ, 459, 185
- Mirabel, I. F., & Rodriguez, L. F. 1994, Nature, 371, 46
- Mirabel, I. F., & Rodriguez, L. F. 1995, Annals NY Academy of Science, 759, 21
- Shakura, N. I., Sunyaev, R. A. 1973, A&A, 24, 337
- Tavani, M., et al. 1996, ApJ, 473, L103
- Tingay, S. J., et al. 1995, Nature, 374, 141




Figure 1: The speeds of propagation of the approaching (solid line) and receding (dashed line) asymmetrical pair of plasmoids calculated from Eqs.(10)-(12) assuming different distances to GRS 1915+105.




Figure 2: The speeds (bottom panel) and the Lorentz factors (top panel) of the bulk motion of the pair of plasmoids calculated for a fixed distance $D = 12.5 \text{ kpc}$ and angular velocities $\mu_{a,r}$ observed from GRS 1915+105 (MR94), but assuming different flux ratios S_a/S_r .